

Table 2.16
Deceased Donor Organ Utilization, 1998 to 2007, Intestine

	Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Total # Donors of Any Organ	5,793	5,824	5,985	6,080	6,190	6,457	7,150	7,593	8,024	8,091
# Intestines Recovered	78	95	87	116	112	123	166	184	185	205
ORPD (1)	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.03
# Intestines Transplanted	68	71	79	112	106	112	146	171	171	197
OTPD (2)	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02
% Transplanted	87.2%	74.7%	90.8%	96.6%	94.6%	91.1%	88.0%	92.9%	92.4%	96.1%
Donor with Brain Death(DBD)										
Total # Donors of Any Organ	5,718	5,737	5,868	5,912	6,001	6,189	6,759	7,029	7,379	7,298
# Intestines Recovered	78	95	87	116	112	122	166	184	185	205
ORPD (1)	0.01	0.02	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03
# Intestines Transplanted	68	71	79	112	106	112	146	171	171	197
OTPD (2)	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.03
% Transplanted	87.2%	74.7%	90.8%	96.6%	94.6%	91.8%	88.0%	92.9%	92.4%	96.1%
Donors After Cardiac Death(DCD)										
Total # Donors of Any Organ	75	87	117	168	189	268	391	564	645	793
# Intestines Recovered	-	-	-	-	-	1	-	-	-	-
ORPD (1)	-	-	-	-	-	0.00	-	-	-	-
# Intestines Transplanted	-	-	-	-	-	0	-	-	-	-
OTPD (2)	-	-	-	-	-	0.00	-	-	-	-
% Transplanted	-	-	-	-	-	0.0%	-	-	-	-
Intestine Donors	78	95	87	116	112	123	166	184	185	205
Intestine Donors (%)	1.3%	1.6%	1.5%	1.9%	1.8%	1.9%	2.3%	2.4%	2.3%	2.5%
Intestine Donors, Other Organs Recovered										
Intestine Only	-	-	1	-	2	-	-	1	-	-
Intestine & Kidney or Pancreas Only	26	32	27	37	39	29	46	46	49	55
Intestine & Liver Only	6	4	5	4	3	5	6	5	2	4

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.

Table 2.16 (Continued)
Deceased Donor Organ Utilization, 1998 to 2007, Intestine

	Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
Intestine Donors, Other Organs Recovered										
Intestine & Thoracic Only	10	9	9	10	7	11	13	13	13	7
Intestine & Other Combinations	36	50	45	65	61	78	101	119	121	139
Intestine Donors, Other Organs Recovered (%)										
Intestine Only	-	-	1.1%	-	1.8%	-	-	0.5%	-	-
Intestine & Kidney or Pancreas Only	33.3%	33.7%	31.0%	31.9%	34.8%	23.6%	27.7%	25.0%	26.5%	26.8%
Intestine & Liver Only	7.7%	4.2%	5.7%	3.4%	2.7%	4.1%	3.6%	2.7%	1.1%	2.0%
Intestine & Thoracic Only	12.8%	9.5%	10.3%	8.6%	6.3%	8.9%	7.8%	7.1%	7.0%	3.4%
Intestine & Other Combinations	46.2%	52.6%	51.7%	56.0%	54.5%	63.4%	60.8%	64.7%	65.4%	67.8%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(1) ORPD: Organs Recovered Per Donor of any organ; (2) OTPD: Organs Transplanted Per Donor of any organ. Organs divided into segments (liver, lung, pancreas, intestine) are counted only once for that organ.

Includes organs recovered for transplant but not transplanted, as well as organs transplanted.

Donors are divided into two mutually exclusive and complete categories. DCDs include patients whose hearts have irreversibly stopped beating, and may also have been known as Non-Heart Beating Donors. DBDs have suffered brain death.